Herpetological Review 36(3): 331; 2005 © 2005 by Society for the Study of Amphibians and Reptiles

BARYCHOLOS PULCHER (NCN). ECUADOR: PROVINCIA DE AZUAY: Tamarindo (79°33'W, 02°47'S, 400 m). 04 August 1996. J.-M. Touzet et al. Universidad San Francisco de Quito & Fund. Herpetológica Orcés, Quito (FHGO-USFQ 623, 748, collected in the leaflitter). Verified by W. Ron Heyer. This endemic frog is known from western lowlands of Ecuador along the transition zone between rainforest and dry seasonal forest from 30 to 600 m; with previous records in the provinces of Esmeraldas, Manabí, Guayas, Los Ríos, and Pichincha (Heyer 1969. Contrib. Sci. Los Angeles Co. Mus. Nat. Hist. 155: 1–14). First province record, extends its range ca. 70 km NE from the nearest known locality (7 km SE Buenavista, province of Oro; Heyer, *op. cit.*).

Submitted by **DIEGO F. CISNEROS-HEREDIA**, College of Biological and Environmental Sciences, Universidad San Francisco de Quito, Ave. Interoceánica y calle Diego de Robles, Campus Cumbayá, Edif. Maxwell. Casilla Postal 17-12-841, Quito, Ecuador; e-mail: diegofrancisco_cisneros@yahoo.com.