

TRACHYBOA BOULENGERI (Northern Eyelash Boa). ECUADOR: PROVINCIA DE ESMERALDAS: Bilsa reserve, Mache Chindul hills (79°45'W, 00°22'S). 7 January 2000. D. Dávalos. Universidad San Francisco de Quito and Fund. Herpetológica Orcés, Quito (FHGO-USFQ 2687-8, 2976); km 17 on Lita-Alto Tambo Road (78°31'W, 00°51'N, 830 m). 14 November 1990. J.-M. Touzet. (FHGO-USFQ 374); km 5 on Lita-Ibarra Road (78°25'W, 00°49'N, 550 m). 18 February 1991. J.-M. Touzet. (FHGO-USFQ 403); Sector Las Golondrinas (79°02'W, 00°20'N, 450 m). 29 November 1994. J.-M. Touzet. (FHGO-USFQ 1121); Zapallo Grande - Río Cayapas (78°57'W, 00°54'N, 90 m). 20 November 1989 – 13 July 1992. J. C. Vieira, J. Coroso, T. Leetz. (FHGO-USFQ 135, 344, 542). Verified by J.-M. Touzet. *Trachyboa boulengeri* occurs along the Chocoan lowlands of Panamá, Colombia and Ecuador. It has been recorded in Ecuador in the provinces of Los Ríos and Pichincha (Perez-Santos and Moreno 1991. *Serpientes de Ecuador*. Monogr. XI. Mus. Reg. Scien. Nat. Torino. 540 pp.). Specimens herein reported represent the westernmost locality of the species and the first record for the province of Esmeraldas, extending its range ca. 90 km NW from the nearest known locality in Pichincha and filling the gap between Colombian and Ecuadorian localities (Perez-Santos and Moreno, *op. cit.*).

Submitted by **DIEGO F. CISNEROS-HEREDIA**, College of Biological and Environmental Sciences, Universidad San Francisco de Quito, Ave. Interoceánica y calle Diego de Robles, Campus Cumbayá, Edif. Maxwell. Casilla Postal 17-12-841, Quito, Ecuador; e-mail: diegofrancisco_cisneros@yahoo.com.